

Mean Sea Surface (MSS) for SWOT

SIO

David Sandwell

Yao Yu

Hugh Harper

CLS/CNES

Philippe Schaeffer

Yannice Faugere

Isabelle Pujol

Gerald Dibarboure

Nicolas Picot

DTU

Ole Andersen

Per Knudsen

Adeli Abulaitijiang

Shengjun Zhang

NOAA

Eric Leuliette

Walter Smith

- Need high resolution MSS for CAL/VAL early in the mission.
- MSS should have long wavelength accuracy from multidecadal repeat-track altimetry (ERM) and short wavelength precision from geodetic mission (GM) phases.
- MSS should vary with time as sea level rises.

MSS(time) for SWOT

MSS(t) = high spatial resolution MSS

+ low spatial resolution MSS (t)

Three groups working on MSS(t) for SWOT:

- 1) CLS constructs MSS2022 = MSS CNES_CLS2021 plus MSS LEADS in Arctic area.
- 2) SIO adds short wavelength MSS from double retracked SSS data.
- 3) DTU develops independent MSS (DTU21), with Arctic focus, and adds lower spatial resolution MSS(t).

MSS(time) for SWOT

Three presentations:

- High spatial resolution MSS (Sandwell – US; Schaeffer – EU)
- Assessment of errors in MSS models (Schaeffer)
- Adding long-wavelength, time - dependent MSS (Andersen)
- Discussion:
 - How important is adding MSS(t)?
 - Would a mean sea surface slope model be useful (along- and cross-track slope)?
 - Does one the MSS need to be embedded in the SWOT product?
 - Will SWOT measure the nadir SSH or the closest reflection SSH?

High Resolution Mean Sea Surface for SWOT

Philippe Schaeffer – CLS

David Sandwell – SIO

Isabelle Pujol – CLS

Ole Andersen – DTU

Improve the shortest wavelengths using High Resolution data

- Goal is to improve wavelengths < 50 km
- Use of GM and drifting data, especially C2 & AltiKa filtered at 5 Hz ($\sim 1,4$ km along track)

Collaborative analysis between CLS & Scripps

HR MSS determination => 2 ways : 2 different dataset and 2 mapping methods are used !

CLS (first step)

Removing oceanic variability



Scripps(second step)

Improving Short wavelengths

Mean Profiles = all ERM Missions
(TP/J1/J2/J3 (& interleave), E2/En/Aka, GFO)

HR Data

Integration of: C2 + AltiKa: **one pass RTK + 5Hz filtering**

S3 => for validation

Observation

SSH – MSLA DUACS

Mapping

Optimal interpolation + noises budget (white & correlated) + noise optimal filtering

Based on CNES_CLS MSS for $\lambda > 100$ Km

HR Data

Integration of: Geosat/J1/J2/En + C2 + AltiKa + S3 : **two-pass RTK + 5 Hz filtering**

Observation

SLOPE combined with HEIGHT

Mapping

Biharmonic splines in tension

SSH – MSLA DUACS : the treatment of the oceanic variability


➤ MSS CLS construction characteristics => **correct each observation for oceanic variability by space-time interpolation of SLA (Map of SLA DUACS = 1 map /day)**

1. Remove (large and meso-scale) seasonal and interannual oceanic variability
2. Remove Sea Level Trend (referenced at an arbitrary time / **mid-1993**)
3. obtain an optimal compromise between mean oceanic content and high-resolution topographic structures

} Provided by Mean Profiles (1Hz)

} Provided by C2 & AltiKa

➤ **goal is to converge towards the “steady state” of the ocean**



▪ DUACS uses the 20y reference period [1993, 2012]

▪ **DUACS convention** : $\langle \text{SLA}_{20y} \rangle_{1993} = 0$

▪ $\langle \text{SLA}_{20y} \rangle_{20y} = \text{Cst value}$ that represents of the MSL increase over the 20y reference period

➤ **Mean SLA over the reference period should be close to 0 (or constant value for DUACS convention) \approx “steady state”**



SSH – MSLA DUACS : the treatment of the oceanic variability

- Mean SLA over the reference period should be close to 0 (or constant value for DUACS convention)

Boxed mean SLA TP/J1/J2 over the MSS reference period [1993, 2012]

SLA TP/J1/j2 [1993, 2012]



TP/J1/J2 mean profile used



Mean of multi-mission DUACS gridded products the MSS reference period [1993, 2012]

gridded SLA DUACS DT-2018[1993, 2012]

MSS CNES15 & MPs used




- nearly cst map as expected => This means that an average calculated over 20 years brings us closer to the steady state !


Rq: mapping method of SLA is limited in high latitude !

SWOT - MSS must be corrected for slope effect

- New MSS will result from a combination of **various altimeter** that are not affected in the same way by the slope of the sea surface.



Slope Correction C2_LRM dH(mm)





[Sandwell and Smith, 2014]

Differences between HR MSS

Scripps_CLS22 (S3) – CNES_CLS22

dH(m)





- These two solutions are very close.
- Differences in amplitude of 1 to 2 cm appear along the geophysical structures that suggest more HR for Scripps model.
- There is a slight impact of ocean variability on areas of strong currents.

Differences between HR MSS

Scripps_CLS22 (S3) – DTU21

dH(m)



- These two solutions remain close in terms of short wavelength content.
- The difference due to the oceanic variability is more significant.

Differences between HR MSS

- Differences are calculated on grids at 1 min resolution.


Bathy > 1000 m

Diff	Nb Points	Mean (cm)	Std (cm) [3σ]
Scripps – CLS	119 439 521	0,06	0,80
CLS - DTU	118 365 843	0,09	1,38
Scripps – DTU	118 861 025	0,02	1,46

Bathy < 1000 m

Diff	Nb Points	Mean (cm)	Std (cm) [3σ]
Scripps – CLS	12 542 354	0,63	3,38
CLS - DTU	12 599 451	0,40	4,99
Scripps – DTU	12 535 188	-0,25	5,22

Pts Involved => Bathy > -1000 m p(1/60)



- The low values of the averages imply that these MSS are "centered" and therefore consistent in term of Sea Level Rise.
- The standard deviation values show that these MSS are close in terms of high-resolution content and also consistent with the expected accuracy of SWOT.
- We note a relative degradation of the accuracy near the coasts which remains one of the major difficulties concerning the processing of altimetric data.
- excluding latitudes higher than 60 degrees gives the same results

Conclusion

- The 3 MSS are "centered in time" and therefore consistent in term of Sea Level Rise.
- In open ocean: MSS are close in terms of high-resolution content and consistent with the expected accuracy of SWOT
- Less consistence near the coast ...

WG MSS recommendation

- The joint use of the SCRIPPS-CLS and DTU MSS for the Cal/Val SWOT activities can allow to decorrelate the imperfections linked to the MSS and to these of SWOT data.

Please look at: Assessment of errors in MSS models presentation for further comparisons